

Measurement of Charmless $B \rightarrow VV$ decays at *BABAR*

Emmanuel Olaiya

Rutherford Appleton Laboratory, Chilton, Didcot, OX11 0QX, UK (for the BABAR Collaboration)

We present results of $B \rightarrow$ vector-vector (VV) and $B \rightarrow$ vector-axial vector (VA) decays $B^0 \rightarrow \phi X$ ($X = \phi, \rho^+$ or ρ^0), $B^+ \rightarrow \phi K^{(*)+}$, $B^0 \rightarrow K^* K^*$, $B^0 \rightarrow \rho^+ b_1^-$ and $B^+ \rightarrow K^{*0} a_1^+$. The largest dataset used for these results is based on $465 \times 10^6 \Upsilon(4S) \rightarrow B\bar{B}$ decays, collected with the *BABAR* detector at the PEP-II B meson factory located at the Stanford Linear Accelerator Center (SLAC).

1. INTRODUCTION

A vector-vector or vector-axial vector B -meson decay can be characterised by three complex helicity amplitudes, A_+ , A_0 and A_- , corresponding to meson helicity states $\lambda = +1, 0$ and -1 , respectively. The A_0 amplitude is expected to dominate due to the $(V - A)$ nature of weak interactions and helicity conservation in strong interactions [1]. Most B decays that arise from tree level $b \rightarrow c$ transitions follow an amplitude heirachy of $|A_0| > |A_+| > |A_-|$. The B meson decays to heavy vector particles with charm such as $B \rightarrow J/\Psi K^*$ show a substantial fraction of the amplitude with transverse polarisation of the mesons ($A_\pm$). This is because the larger the meson mass, the weaker the amplitude heirachy. Therefore the amplitude heirachy $|A_0| > |A_+| > |A_-|$ is expected to be more significant in B decays to light vector particles in both penguin [2–4] and tree-level [1] transitions. In the tree-level $b \rightarrow u$ transitions, such as $B^0 \rightarrow \rho^+ \rho^-$ [5], $B^+ \rightarrow \rho^0 \rho^+$ [6], and $B^+ \rightarrow \omega \rho^+$ [7], the dominance of longitudinal polarisation has been confirmed by the *BABAR* and Belle experiments. However a large fraction of transverse polarisation was first observed by *BABAR* and confirmed by Belle for the decays $B \rightarrow \phi K^*$ and $B \rightarrow \rho K^*$, which is a significant departure from the expected dominance of the longitudinal amplitude.

This polarisation anomaly in the vector-vector decays $B \rightarrow \phi K^*$ and $B \rightarrow \rho K^*$ suggests other contributions to the amplitude, that were previously neglected. This possibility has generated great interest in vector-vector decays, motivating a number of proposed contributions from physics beyond the standard model [8]. In addition there are mechanisms within the standard model which have been proposed to explain the anomaly such as annihilation penguin [4, 9] or electroweak penguin, or QCD rescattering [10]. The key to understanding this anomaly could be obtained by studying other vector-vector and vector-axial vector decays such as $B \rightarrow \phi V$, $B \rightarrow K^* K^*$, $B \rightarrow \rho b_1$, $B \rightarrow K^* a_1$ and comparing experimental results with theory.

2. METHOD

The angular distribution of the B decay products can be expressed as a function of three helicity angles which describe the alignment of the particles in the decay chain. For two-body decays the polarisation is normally chosen as the direction of the daughters in the center of mass of the parent [11], and for three-body decays the normal to the decay plane [12]. For $B \rightarrow VV$ and $B \rightarrow VA$ decays the angular distribution in the helicity frame is given by the equation:

$$\frac{d^2\Gamma}{d\cos\theta_1 d\cos\theta_2} = \frac{9}{4}[f_L \cos^2\theta_1 \cos^2\theta_2 + \frac{1}{4}(1 - f_L) \sin^2\theta_1 \sin^2\theta_2] \quad (1)$$

where $\theta_{1,2}$ are the helicity angles and f_L is the fraction of longitudinal polarisation and is equal to $|A_0|^2/(|A_0|^2 + |A_{-1}|^2 + |A_{+1}|^2)$. Here we have integrated over the azimuthal angle ϕ (shown in Figure 1). We calculate the fraction of longitudinal polarisation f_L by fitting the angular distributions θ_1 and θ_2 .




Figure 1: Definition of helicity angles using the decay $B \rightarrow \rho^+ b_1$. The angle θ_1 corresponds to b_1 and θ_2 corresponds to ρ^+ .

3. RESULTS

3.1. $B^0 \rightarrow \phi\phi$, $B^+ \rightarrow \phi\rho^+$ and $B^0 \rightarrow \phi\rho^0$

The decay $B^0 \rightarrow \phi\phi$ is an OZI suppressed process with an expected branching fraction in the range $(0.1 \text{ to } 3.0) \times 10^{-8}$ in the standard model [13–15]. The decays $B^+ \rightarrow \phi\rho^+$ and $B^0 \rightarrow \phi\rho^0$ are pure $b \rightarrow d$ loop processes with expected branching fractions in the range $(0.2 \text{ to } 5.3) \times 10^{-7}$ [16–24].

Using a data sample of 384×10^6 $B\bar{B}$ pairs we observed no signal for these decays, consistent with standard model expectations. We obtain a 90% confidence level upper limit for the decays $B^0 \rightarrow \phi\phi$, $B^+ \rightarrow \phi\rho^+$ and $B^0 \rightarrow \phi\rho^0$ of 2×10^{-7} , 30×10^{-7} and 3.3×10^{-7} [25], respectively.

3.2. $B^+ \rightarrow \phi K^{(*)+}$

We search for $B^+ \rightarrow \phi K^{(*)+}$ decays using three final states of the $K^{(*)+}$ decay: $K_S^0\pi^+$, $K^+\pi^0$ and $K^+\pi^+\pi^-$, where $K_S^0 \rightarrow \pi^+\pi^-$ and $\pi^0 \rightarrow \gamma\gamma$. Previous studies have been limited to the two body $K_J^* \rightarrow K\pi$ decays, thus considering only the spin-parity K_J^* states with $P = (-1)^J$. We now consider three-body states $K_J^* \rightarrow K\pi\pi$ which include $P = (-1)^{J+1}$ states such as K_1 and K_2 . Using a data sample of 465×10^6 $B\bar{B}$ pairs we perform an amplitude analysis to measure branching fractions and polarisations for $B \rightarrow \phi K^*$ decays [26]. The results of the amplitude analysis are shown in Table I.

3.3. $B^0 \rightarrow K^{*0}\bar{K}^{*0}$ and $B^0 \rightarrow K^{*+}K^{*-}$

Decays proceeding via electroweak and gluonic $b \rightarrow d$ penguin diagrams have been measured in the decays $B \rightarrow \rho\gamma$ [27] and $B^0 \rightarrow K^0\bar{K}^0$ [28]. However, the decay $B^0 \rightarrow K^{*0}\bar{K}^{*0}$ proceeds via both electroweak and gluonic $b \rightarrow d$ penguin loops to two vector particles. Theoretical calculations for $B^0 \rightarrow K^{*0}\bar{K}^{*0}$ branching fractions cover a range $(0.16\text{--}0.96) \times 10^{-6}$ [19, 21, 29]. The latest predictions by Beneke *et al.* [15] estimate a branching fraction of $(0.6^{+0.1+0.3}_{-0.1-0.2}) \times 10^{-6}$ and $f_L = 0.69 \pm 0.01^{+0.16}_{-0.20}$. Using a data sample of 383×10^6 $B\bar{B}$ pairs we observe the decay $B^0 \rightarrow K^{*0}\bar{K}^{*0}$ for the first time, measuring a branching fraction of $(1.28^{+0.35}_{-0.30} \pm 0.11) \times 10^{-6}$ and $f_L = 0.80^{+0.10}_{-0.12} \pm 0.06$ [30], which is consistent with theoretical predictions.

The decay $B^0 \rightarrow K^{*+}K^{*-}$ is expected to occur through a $b \rightarrow u$ quark transition via a W -exchange, and consequently have a small branching fraction with predictions of $(0.09^{+0.05+0.12}_{-0.03-0.10}) \times 10^{-6}$ [15] and $(0.1 \pm 0.0 \pm 0.1) \times 10^{-6}$ [2] both based on QCD factorisation. Using a data sample of 454×10^6 $B\bar{B}$ pairs we observe no signal and obtain a 90% confidence upper limit of 2.0×10^{-6} [31] which is two orders of magnitude more sensitive than the previous measurement [32].

Table I: Number of signal events, significance, longitudinal polarisation fraction and branching fraction measurements from the $B^+ \rightarrow \phi K^{(*)+}$ amplitude analysis.

Mode	n_{sig}	s (σ)	f_L	B (10^{-6})
$\phi K_1(1270)^+$	$116 \pm 26 \pm 14$	5.0	$0.46^{+0.12+0.03}_{-0.13-0.07}$	$6.1 \pm 1.6 \pm 1.1$
$\phi K_1(1400)^+$	$7 \pm 39 \pm 17$	0.2		< 3.2 ($0.3 \pm 1.6 \pm 0.7$)
$\phi K_1(1430)^+$		5.5	$0.80^{+0.09}_{-0.10} \pm 0.03$	$(8.4 \pm 1.8 \pm 0.9)$
$\rightarrow K_S^0 \pi^+$	$27 \pm 6 \pm 3$			
$\rightarrow K^+ \pi^0$	$39 \pm 8 \pm 4$			
$\rightarrow K^+ \pi^+ \pi^-$	$64 \pm 14 \pm 6$			
$\phi(K\pi)_0^{*+}$		8.2		$(8.3 \pm 1.4 \pm 0.8)$
$\rightarrow K_S^0 \pi^+$	$48 \pm 8 \pm 4$			
$\rightarrow K^+ \pi^0$	$80 \pm 13 \pm 7$			
$\phi K_0^*(1430)^+$				$(7.0 \pm 1.3 \pm 0.9)$
$\phi K^*(1410)^+$	$61 \pm 31^{+11}_{-31}$	< 2.0		< 4.8 ($2.4 \pm 1.2^{+0.4}_{-1.2}$)
$\phi K^*(1770)^+$	$90 \pm 32^{+39}_{-46}$	< 2.0		< 16.0
$\phi K^*(1820)^+$	$122 \pm 40^{+20}_{-83}$	< 2.0		< 23.4

3.4. $B^0 \rightarrow \rho^+ b_1^-$

Cheng and Yang [2] have extended calculations by Beneke *et al.*[15] to include vector-axial vector (VA) decays. They predict a $B^0 \rightarrow \rho^+ b_1^-$ branching fraction of $(32.7^{+16.5+12.1}_{-14.7-4.7}) \times 10^{-6}$. This measurement combines the branching fractions of both $B^0 \rightarrow \rho^+ b_1^-$ and $B^0 \rightarrow \rho^- b_1^+$, noting that the latter decay is suppressed by G -parity conservation. They also predict the longitudinal polarisation fraction to be $f_L = 0.96^{+0.01}_{-0.02}$.

Using a data sample of 465×10^6 $B\bar{B}$ pairs we find no evidence of a signal and obtain a 90% confidence level upper limit of 1.7×10^{-6} [33].

3.5. $B^+ \rightarrow K^{*0} a_1^+$

Recent searches of charmless decays with axial-vectors have revealed comparatively large branching fractions compared to other charmless decays, in the range $(15 - 35) \times 10^{-6}$ [34]. Available $B^+ \rightarrow K^{*0} a_1^+$ theoretical estimates are based on naïve factorisation with an expected branching fraction of 0.51×10^{-6} [35] and on QCD factorisation with an expected branching fraction of $(9.7^{+4.9+32.9}_{-3.5-2.4}) \times 10^{-6}$ and a longitudinal polarisation fraction f_L of $0.38^{+0.51}_{-0.40}$ [2].

Using a data sample of 465×10^6 $B\bar{B}$ pairs we find no evidence for signal and measure a branching fraction of $(1.4^{+0.8+1.4}_{-1.0-1.4}) \times 10^{-6}$ corresponding to a 90% confidence level upper limit of 3.2×10^{-6} [36].

4. CONCLUSION

Using larger datasets, the *BABAR* experiment has provided more precise $B \rightarrow VV$ measurements, further supporting the smaller than expected longitudinal polarisation fraction of $B \rightarrow \phi K^*$. Additional B meson to vector-vector and vector-axial vector decays have also been studied with a view to shedding light on the polarisation anomaly. Taking into account the available errors, we find no disagreement between theory and experiment for these additional decays.

Acknowledgments

I would like to thank the organisers of the ICHEP 08 conference, the *BABAR* collaboration and the UK Science and Technology Facilities Council (STFC).

References

- [1] A. Ali *et al.*, Z. Physik **C1**, 269 (1979);
M. Suzuki, Phys. Rev. **D64**, 117503 (2001).
- [2] H. Y. Cheng and K. C. Yang, Phys. Lett. **B511**, 40 (2001).
- [3] C. H. Chen, Y. Y. Keum and H. n. Li, Phys. Rev. **D66**, 054013 (2002).
- [4] A. L. Kagan, Phys. Lett **601**, 151 (2004);
Y. Grossman, Int. J. Mod. Phys. **A19**, 907 (2004).
- [5] Belle Collaboration, A. Somov *et al.*, Phys. Rev. Lett. **96**, 171801 (2006);
BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. **D76**, 052007 (2007).
- [6] Belle Collaboration, J. Zhang *et al.*, Phys. Rev. Lett. **91**, 221801 (2003);
BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. Lett **97**, 261801 (2006).
- [7] BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. **D74**, 051102 (2006).
- [8] E. Alvarez *et al.*, Phys. Rev. D **70**, 115014 (2004); P. K. Das and K. C. Yang, Phys. Rev. D **71**, 094002 (2005);
C. H. Chen and C. Q. Geng, Phys. Rev. D **71**, 115004 (2005).
- [9] H. n. Li and S. Mishima, Phys. Rev. D **71**, 054025 (2005); C.-H. Chen *et al.*, Phys. Rev. D **72**, 054011 (2005);
M. Beneke *et al.*, Phys. Rev. Lett. **96**, 141801 (2006), arXiv:hep-ph/0612290; C.-H. Chen and C.-Q Geng, Phys.
Rev. D **75**, 054010 (2007); A. Datta *et al.*, arXiv:0705.3915[hep-ph].
- [10] C. W. Bauer *et al.*, Phys Rev. D **70**, 054015 (2004); P. Colangelo *et al.*, Phys. Lett. B **597**, 291 (2004).
- [11] M. Jacob and G. C. Wick, Ann. Phys. **7**, 404 (1959).
- [12] S. M. Berman and M. Jacob, Phys. Rev. **139**, 1023 (1965).
- [13] S. Bar-Shalom, G. Eilam, and Y. D. Yang, Phys. Rev. D **67**, 014007 (2003).
- [14] C. D. Lu *et al.*, Eur. Phys. Jour. C **41**, 311 (2005).
- [15] M. Beneke, J. Rohrer and D. Yang, Nucl. Phys. B **774**, 64 (2007).
- [16] D. Du and Z. Xing, Phys. Lett. B **312**, 199 (1993).
- [17] M. Gronau and J. L. Rosner, arXiv:0806.3584.
- [18] A. Deandrea *et al.*, Phys. Lett. B **320**, 170 (1994).
- [19] A. Ali *et al.*, Phys. Rev D **58**, 094009 (1998).
- [20] Y. H. Chen *et al.*, Phys. Rev. D **60**, 094014 (1999).
- [21] W. Zou and Z. Xiao, Phys. Rev. **D72**, 094026 (2005).
- [22] C. D. Lu *et al.*, Chin. Phys. Lett. **23**, 2684 (2006).
- [23] J. Li. *et al.* hep-ph/0607249.
- [24] S. Bao, F. S. Su, Y.-L Wu, and C.Zhuang, arXiv:0801.2596.
- [25] B. Aubert *et al.* (BABAR Collaboration), arXiv:0807.3935 [accepted for publication in PRL].
- [26] B. Aubert *et al.* (BABAR Collaboration), arXiv:0806.4419 [accepted for publication in PRL].
- [27] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. Lett. **98**, 151802 (2007); D. Mohapatra *et al.* (Belle Collaboration), Phys. Rev. Lett. **96**, 221601 (2006).
- [28] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. Lett. **97**, 171805 (2006); S.-W. Lin *et al.* (Belle Collaboration) Phys. Rev. Lett. **98**, 181804 (2007).
- [29] L. Chau *et al.*, Phys. Rev. D **45**, 3143 (1992).
- [30] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. Lett. **100**, 081801 (2008).
- [31] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. D **78**, 051103 (2008).
- [32] R. Godang *et al.* (CLEO Collaboration), Phys. Rev. Lett. **88**, 021802 (2002).
- [33] W.T.Ford, “Hadronic B_u and B_d decays”, arXiv:0810.0494.
- [34] BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. Lett **99**, 261801 (2007); B. Aubert *et al.*, Phys. Rev. Lett **100**, 051803 (2008).
- [35] G. Calderón, J. H. Muñoz and C. E. Vera, Phys. Rev. D **76**, 094019 (2007).
- [36] B. Aubert *et al.* (BABAR Collaboration), arXiv:0808.0579.